

Prospect of creating a composite fermi/bose superfluid

Eddy Timmermans,¹ Kyoko Furuya,² Peter W. Milonni,¹
and Arthur K. Kerman³

¹ Theoretical Division (T-4), Los Alamos National Laboratory, Los Alamos, NM 87545

² Universidade Estadual de Campinas, Instituto de Física ‘Gleb Wataghin’
c.p. 6165, 13083-970, Campinas, SP, Brazil

³ Center for Theoretical Physics, Laboratory for Nuclear Science and
Department of Physics, Massachusetts Institute of Technology,
Cambridge, MA 02139

Abstract

We show that composite fermi/bose superfluids can be created in cold-atom traps by employing a Feshbach resonance or coherent photoassociation. The bosonic molecular condensate created in this way implies a new fermion pairing mechanism associated with the exchange of fermion pairs between the molecular condensate and an atomic fermion superfluid. We predict macroscopically coherent, Josephson-like oscillations of the atomic and molecular populations in response to a sudden change of the molecular energy, and suggest that these oscillations will provide an experimental signature of the pairing.

PACS numbers(s):03.75.Fi, 05.30.Jp, 32.80Pj, 67.90.+z

Typeset using REVTeX

The cold-atom technology that has led to the observation of dilute-gas Bose-Einstein condensates (BEC's) [1] enables the creation and study of novel superfluids having a degree of flexibility that is highly unusual in traditional low temperature physics. In addition to the observed boson gas superfluids, the prospect of studying fermion superfluidity in the gas phase provides a powerful motivation [2] for the atom cooling efforts with fermion isotopes [3]. S-wave Cooper pairing – the most likely mechanism for achieving fermion superfluidity in the low-density gases – relies on the mutual attraction of two distinguishable types of fermions [4]. In this Letter, we show that some of the techniques that are being developed [5] to increase the mutual attraction and, hence, the critical temperature for Cooper-pairing, create a fundamentally novel superfluid state. Specifically, the interaction-altering schemes that involve a coherent atom-molecule coupling such as the Feshbach resonance [6] and coherent photoassociation [7] create a condensate of bosonic molecules mutually coherent with the atomic Cooper-paired gas. In treating the dynamics of this system we note an interesting experimental signature: the system responds to a sudden change of the single-molecule energy by means of macroscopically coherent, Josephson-like oscillations of the atomic and molecular populations.

Here we discuss the effects on a mixture of ± 1 fermion atoms of a Feshbach resonance that coherently converts $\pm$ atom pairs to molecules m . Restricting ourselves to the relevant s-wave processes, the coupling is described by the term

$$H_{a-m} = \int d^3\mathbf{r} \ \alpha \left[\hat{\psi}_m^\dagger \hat{\psi}_{+1} \hat{\psi}_{-1} + \text{h.c.} \right] \quad (1)$$

in the Hamiltonian. Earlier [6], we had pointed out that such interaction can produce mutually coherent atom-molecule boson superfluids. In the fermion gas mixture, we now show that this interaction coherently couples an atomic paired fermion superfluid to a BEC of m -molecules. We characterize boson superfluidity in the broken symmetry description by the appearance of a complex-valued order parameter, the

condensate field ϕ : $\phi(\mathbf{r}) = \langle \hat{\psi}(\mathbf{r}) \rangle$, where $\hat{\psi}$ is the bosonic annihilation field operator and $\mathbf{r}$ the position. Similarly, we characterize s-wave pairing of fermionic atoms in internal states ± 1 by a pairing-field order parameter $\langle \hat{\psi}_\sigma(\mathbf{r}) \hat{\psi}_{-\sigma}(\mathbf{r}) \rangle = F_\sigma(\mathbf{r}) \neq 0$, where $\sigma = \pm 1$. It is not difficult to see that the interaction (1) implies the presence of a bose condensate of molecules ($\langle \hat{\psi}_m \rangle \neq 0$) if the atoms are fermi-paired ($\langle \hat{\psi}_{+1} \hat{\psi}_{-1} \rangle \neq 0$). The equation of motion for $\langle \hat{\psi}_m \rangle = \phi_m$, obtained by taking the expectation value of the Heisenberg equation $i\hbar \partial \hat{\psi}_m / \partial t = [\hat{\psi}_m, \hat{H}]$, contains a source term $\alpha \langle \hat{\psi}_{+1} \hat{\psi}_{-1} \rangle \neq 0$, implying that ϕ_m cannot remain zero. Moreover, as we will show, the interaction (1) plays a more fundamental role than in the BEC analogue: the atom-molecule interaction *causes* the superfluidity and provides a new pairing mechanism.

From a theoretical point of view, however, this mechanism is not entirely new as the Hamiltonian (1) has been proposed as a phenomenological model for high T_c superconductivity [8]. However, experimental investigations have cast serious doubt on the validity of the so-called fermion-boson model to explain high T_c superconductivity. It is then important to realize that, in contrast to the phenomenological description of high T_c superconductors, the atomic gases are described by a first principle's theory with parameters that are calculated and measured unambiguously. We also note that the trapped atom systems have a different and much wider range of experimentally controllable variables: the external potentials, inter-particle scattering lengths, fermion densities, relative numbers of unlike fermions, the strength (α) of the atom-molecule coupling and the value of the single molecule energy can all be varied. In particular, the real time variation of the single molecule energy suggests testing the mutual long-range coherence by observing Josephson-like oscillations in the atomic and molecular populations. Prompted by this exciting possibility we develop a time-dependent mean-field description of the dynamics, different from the

usual statistical treatments of the fermion-boson model.

We start with the Heisenberg equations for the field operators, $\hat{\psi}_m$ and $\hat{\psi}_{\pm 1}$. As we will justify a posteriori, the scattering length interactions can be neglected and we obtain

$$\begin{aligned} i\hbar\dot{\hat{\psi}}_{+1} &= \left[-\frac{\hbar^2\nabla^2}{2m} + V_{+1} + \epsilon_{+1} \right] \hat{\psi}_{+1} - \alpha\hat{\psi}_m\hat{\psi}_{-1}^\dagger, \\ i\hbar\dot{\hat{\psi}}_{-1} &= \left[-\frac{\hbar^2\nabla^2}{2m} + V_{-1} + \epsilon_{-1} \right] \hat{\psi}_{-1} + \alpha\hat{\psi}_m\hat{\psi}_{+1}^\dagger, \\ i\hbar\dot{\hat{\psi}}_m &= \left[-\frac{\hbar^2\nabla^2}{4m} + V_m + \epsilon_m \right] \hat{\psi}_m + \alpha\hat{\psi}_{+1}\hat{\psi}_{-1}, \end{aligned} \quad (2)$$

where $V_{\pm 1}$ and V_m denote the external potentials and ϵ_m and $\epsilon_{\pm 1}$ the internal energies. The fact that fermion atoms of different spin projection can have significantly different energies in the external field of magnetic traps (e.g., $\epsilon_{-1} \ll \epsilon_{+1}$) does not necessarily prevent their pairing. Though the conversion of atoms from the $+1$ to the -1 state would lower the energy, the necessary spin-flip interaction may be negligible, as demonstrated by the simultaneous trapping of essentially stable populations of atoms in non-degenerate internal states [9]. Such systems reach a quasi-equilibrium, characterized, in this case, by fixed fermion numbers: a total of N_+ ($+1$)–atoms and N_- (-1)–atoms, with respective chemical potentials μ_+ and μ_- . Chemical equilibrium further dictates that $\mu_m = \mu_+ + \mu_-$, where μ_m is the molecular chemical potential. A simple transformation of the field operators, $\hat{\psi}'_\sigma = \hat{\psi}_\sigma \exp \left[(i/\hbar) \int^t \{ \epsilon_\sigma(\tau) + \mu_\sigma \} d\tau \right]$ and $\hat{\psi}'_m = \hat{\psi}_m \exp \left[(i/\hbar) \int^t \{ \epsilon_+(\tau) + \epsilon_-(\tau) + \mu_m \} d\tau \right]$, illustrates that the dynamics does not depend on the fermion energy difference, $\epsilon_+ - \epsilon_-$, but only on the molecular excitation energy $\epsilon = \epsilon_m - [\epsilon_+ + \epsilon_-]$. Indeed, the transformed operators satisfy

$$\begin{aligned} i\hbar\dot{\hat{\psi}}'_\sigma &= \hat{h}_\sigma\hat{\psi}'_\sigma - \sigma\alpha\hat{\psi}'_m\hat{\psi}'_{-\sigma}^\dagger, \\ i\hbar\dot{\hat{\psi}}'_m &= \hat{h}_m\hat{\psi}'_m + \alpha\hat{\psi}'_+\hat{\psi}'_- , \end{aligned} \quad (3)$$

where we drop the primes and define $\hat{h}_\sigma = \left[-\frac{\hbar^2\nabla^2}{2m} + V_\sigma - \mu_\sigma \right]$, $\hat{h}_m =$

$\left[-\frac{\hbar^2 \nabla^2}{4m} + V_m + \epsilon - \mu_m\right]$. As in Cooper-pairing [10], the annihilation of a fermion is accompanied by the creation of a fermion of opposite “spin” [11]. Accordingly, the quasiparticle state of quantum number j and spin σ , $|j, \sigma\rangle$, is a mixture of a σ particle and a $-\sigma$ hole. We characterize the fermion pairing field by the particle and hole amplitudes for bringing the many-body state $|0\rangle$ to the excited $|j, \sigma\rangle$ -state: $u_{j,\sigma}(\mathbf{r}, t) = \langle 0 | \hat{\psi}_\sigma(\mathbf{r}, t) | j, \sigma \rangle$, $v_{j,\sigma}(\mathbf{r}, t) = -\sigma \langle 0 | \hat{\psi}_\sigma^\dagger(\mathbf{r}, t) | j, -\sigma \rangle$. We obtain the zero temperature equations of motion by taking the excitation amplitudes of the fermion Heisenberg equations and the expectation value of the $\dot{\hat{\psi}}_m$ equation. In the assumption of molecular condensate coherence, $\langle \hat{\psi}_m \rangle \approx \phi_m$, and $\langle 0 | \hat{\psi}_m^\dagger \hat{\psi}_\sigma | j, \sigma \rangle \approx \phi_m^* u_{j,\sigma}$ [12], this mean-field procedure gives Bogoliubov de-Gennes (BDG)-like equations for the u/v -parameters coupled to a Gross-Pitaevskii (GP)-like equation for ϕ_m :

$$\begin{aligned} i\hbar \frac{d}{dt} \begin{pmatrix} u_{j,\sigma} \\ v_{j,-\sigma} \end{pmatrix} &= \begin{pmatrix} \hat{h}_\sigma & -\alpha\phi_m \\ -\alpha\phi_m^* & -\hat{h}_{-\sigma} \end{pmatrix} \begin{pmatrix} u_{j,\sigma} \\ v_{j,-\sigma} \end{pmatrix}, \quad \sigma = \pm 1, \\ i\hbar \frac{\partial \phi_m}{\partial t} &= \hat{h}_m \phi_m + \alpha F_{+1}, \end{aligned} \quad (4)$$

where $F_{+1} = \langle \hat{\psi}_{+1} \hat{\psi}_{-1} \rangle = \sum_j \langle 0 | \hat{\psi}_{+1} | j, +1 \rangle \langle j, +1 | \hat{\psi}_{-1} | 0 \rangle = \sum_j u_{j,+} v_{j,-}^*$ represents the pairing field. The anticommutator relations satisfied by the $\hat{\psi}_\sigma$ -fields imply that $F_{+1} = -F_{-1}$, and that $\sum_j [u_{j,\sigma}(\mathbf{r}) u_{j,\sigma}^*(\mathbf{r}') + v_{j,\sigma}(\mathbf{r}) v_{j,\sigma}^*(\mathbf{r}')] = \delta(\mathbf{r} - \mathbf{r}')$.

In a homogeneous system, $V_+ = V_- = 0$, contained in a macroscopic volume Ω , the excited many-body states have definite momentum, $|j, \sigma\rangle \rightarrow |\mathbf{k}, \sigma\rangle$ and the $u_{j,\sigma}, v_{j,\sigma}$ mode functions are plane waves, $u_{j,\sigma}(\mathbf{r}, t) = u_{k,\sigma}(t) \exp(i\mathbf{k} \cdot \mathbf{r})/\sqrt{\Omega}$. Fermion anticommutation implies $|u_{k,\sigma}|^2 + |v_{k,\sigma}|^2 = 1$, and the $\hat{h}_\sigma$ -operators of (4) give real numbers, $\hat{h}_\sigma \rightarrow h_{k,\sigma} = [\hbar^2 k^2/2m - \mu_\sigma]$. For $N_+ = N_- = N$, the ideal situation to achieve superfluidity, $\mu_+ = \mu_- = \mu$, the $\sigma = \pm 1$ equations are identical,

$$\begin{aligned} i\hbar \dot{u}_k &= h_k u_k - \alpha \phi_m v_k \\ i\hbar \dot{v}_k &= -\alpha \phi_m^* u_k - h_k v_k \end{aligned}$$

$$i\hbar\dot{\phi}_m = (\epsilon - 2\mu)\phi_m + \alpha F_+ , \quad (5)$$

where $F_+ = \sum_{\mathbf{k}} u_k v_k^* / \Omega$ and we have dropped the spin subscripts. The single molecule energy, ϵ , can be varied experimentally by altering the magnetic field strength in the Feshbach resonance, or the laser detuning in the case of coherent photoassociation.

Statics In the static ($\dot{\phi}_m = 0$) off-resonant limit where ϵ greatly exceeds all other single particle energies including the fermi-energy $\epsilon_F = \hbar^2 k_F^2 / 2m$, where k_F denotes the Fermi-momentum, $k_F = (6\pi^2 n)^{1/3}$ and $n = N/\Omega$, few molecules are formed and $\mu \approx \epsilon_F$. We recover the BDG equation from (4) by substituting ϕ_m from the GP-equation, $\phi_m = -(\alpha F_{+1})/(\epsilon - 2\mu)$, in the u/v -equations which gives gives the usual Fermi-pairing equations [10] with an effective scattering length

$$a_{\text{eff}} = - \left(\frac{m}{4\pi\hbar^2} \right) \frac{\alpha^2}{\epsilon - 2\mu} , \quad (6)$$

similar to the effective scattering length of the binary atom Feshbach resonance [6]. Not surprisingly, the off-resonant critical temperature T_c obtained in the finite-temperature mean-field treatment takes on the form of a negative scattering length T_c : $T_c \approx T_{c,eff} = 0.6 \exp(-\pi/2k_F |a_{eff}|)$. Near-resonance, T_c follows from coupled integral equations [8], although our numerical results for a realistic system (a Rb_{86} -mixture at $n \approx 2. \times 10^{13} \text{ cm}^{-3}$, $\alpha\sqrt{n} = \epsilon_F$) differed from $T_{c,eff}$ by less than a factor of 2 (giving near-resonant T_c 's $\sim 10^{-7}$ K) [13].

The static u and v -amplitudes in (5) evolve according to the excitation energies $E(k)$ of the $|\mathbf{k}, \sigma\rangle$ -state, $u_k(t) = u_k \exp[-iE(k)t/\hbar]$, $v_k(t) = v_k \exp[-iE(k)t/\hbar]$. The resulting u/v -equations have a non-trivial solution if

$$E(k) = \sqrt{\left(\hbar^2 k^2 / 2m - \mu \right)^2 + |\alpha \phi_m|^2} , \quad (7)$$

giving a superconductor-like dispersion with gap $|\alpha \phi_m|$. Since the gap measures the success of the fermion pairing in lowering the overall energy, Eq. (7) illustrates that it

is the mutual coherence of the Bose and Fermi fields that pairs the fermions, i.e. the boson and fermion fields maintain macroscopic coherence by ‘feeding off’ each other.

The static u and v ‘coherence’ factors that follow from (5), $u_k^2 = [1 + h_k/E(k)]/2$ and $v_k^2 = [1 - h_k/E(k)]/2$, characterize the fermion pairing physics. The occupation number of the single fermion $\mathbf{k}$ -state is given by v_k^2 and the Fourier-transform of $g(k) = v_k/u_k$ gives the wavefunction $\varphi(\mathbf{r} = \mathbf{r}_i - \mathbf{r}_j)$ of the Cooper pairs. The range of the pair wavefunction, φ , $L_c^2 = \int d^3r r^2 \varphi(\mathbf{r}) / \int d^3r \varphi(\mathbf{r})$, follows from the long wavelength behavior of $g(k) \approx g(0)[1 - (kL_c)^2/6]$, $k \rightarrow 0$. We find that $\hbar^2/(2mL_c^2) = E(k=0)/6 = \sqrt{4\mu^2 + (\alpha\phi_m)^2}/6$ and L_c is largest in the off-resonant limit where $\mu \approx \epsilon_F$ and $|\alpha\phi_m| \ll \epsilon_F$, so that $L_c \approx \sqrt{3}k_F^{-1}$. Typically [14], the local value of k_F^{-1} exceeds the length scale on which the density of a trapped degenerate fermion system varies so that the homogeneous results also apply to the trap system in the sense of a Thomas-Fermi description.

Provided $|\alpha\phi_m| \ll \mu$, we can approximate the F_+ -integral analytically and the properly renormalized [15] GP-like equation yields $\phi_m \approx 8e^{-2}(\mu/\alpha) \exp\left(-\frac{\pi}{2k_F|a_{eff}|}\right)$, where $\mu \approx \epsilon_F$. From the exponential dependence on $(k_F a_{eff})^{-1}$ (as compared to $\phi_m \propto \epsilon^{-1}$ in the resonant BEC [6]) it follows that only close to resonance does a measurable molecular field appear, $|a_{eff}| \geq 0.4k_F^{-1}$. This regime is accessible to Feshbach resonance experiments. In that case, the limitation in $\epsilon(a_{eff})$ corresponds to a magnetic field interval around the resonant value B_0 , $|\delta B| < \delta\overline{B}$, where $\delta B = B - B_0$. We determine the interval size $\delta\overline{B}$ from $|a_{eff}| \geq 0.4 k_F^{-1}$ with [16] $a_{eff} = -a_s(\Delta/\delta B)$, where a_s denotes the ordinary scattering length of the unlike fermion interaction and Δ represents the measured magnetic field ‘width’: $\delta\overline{B} = 2.5 \times (|a_0|k_F)|\Delta|$. To obtain representative numbers, we consider the Na_{23} resonances observed by Stenger et al. [16] for which $|\Delta| = 0.1G, 1.G$ and $4G$. Taking $n = 10^{12}cm^{-3}$, [4] and $a_s = 3.3 nm$ (the Na triplet scattering length), we find $\delta\overline{B} = 0.03 |\Delta| = 3mG, 30mG, 0.12G$ exceeding

current experimental magnetic field resolution (typically better than $1mG$). Also, within $|\delta B| \leq \delta \bar{B}$, $|a_{eff}| = |\Delta/\delta B|a_s > 30$. a_s , which justifies our approximation of neglecting the a_s -interactions in (2). As the system is tuned closer to resonance, ϕ_m and μ need to be solved self-consistently from (5) and the condition of particle conservation: $N = \Omega|\phi_m|^2 + \sum_k v_k^2$.

Dynamics As mutually coherent superfluids, the atom-molecule system responds to a sudden change of α or ϵ by means of macroscopically coherent Josephson-like atomic and molecular population oscillations. In the figure, we show the molecular condensate population as ϵ varies in time [17]: at $t=\hbar/\epsilon_F$, ϵ was suddenly raised from $8\epsilon_F$ to $14\epsilon_F$, and returned to its initial value at $t = 1.2\hbar/\epsilon_F$. The time-unit $\hbar/\epsilon_F = 2.1 A(n/n_0)^{-2/3}\mu\text{sec}$, where A is the atomic number and n_0 denotes a reference density of realistic value, $n_0 = 10^{12}cm^{-3}$, indicates time scales that are readily accessible to the available technology. Note that a population oscillation in a binary atom collision would start when the atoms meet, which happens ‘randomly’ in a classical description of the atomic center-of-mass motion, thereby washing out any such oscillations in a many-body average. Thus, macroscopically coherent atom-molecule oscillations constitute a genuine manifestation of mutually coherent long-range order.

Given the problems encountered in the evaporative cooling of neutral fermion atoms [3], it may be of future interest to use coherent molecule formation in an alternative cooling scheme, such as the example we now discuss. (i) We photoassociate finite temperature atoms to molecules. (ii) We switch off the lasers and cool the bosonic molecules evaporatively to BEC. (iii) We turn the lasers on and sweep the detuning (which plays the role of ϵ) of the resulting fermi/bose superfluid from negative to positive values. If we can neglect spontaneous photon emission and other heating and loss-processes, this procedure results in an ultra-cold fermi/bose system of mostly fermionic atoms. (iv) We switch off the lasers.

In conclusion, we have pointed out that the coherent atom-molecule interactions available in present-day atom cooling experiments suggest the exciting prospect of creating a cold atom superfluid of novel structure, consisting of mutually coherent field components of different spin statistics. Although this novel type of fermion pairing mechanism was previously proposed as a phenomenological model for high T_c superconductivity, the atom trap systems could provide the first realization. In addition to the advantage of being described by a well-defined, first principles Hamiltonian, we show that the added flexibility of controlling the detuning can be used to diagnose this new long-range coherence.

REFERENCES

- [1] M. H. Anderson, et al., *Science* **269**, 198 (1995); C. C. Bradley, et al., *Phys. Rev. Lett.* **75**, 1687 (1995); K. B. Davis, et al., *Phys. Rev. Lett.* **75**, 3969 (1995); W. Ketterle, *Physica B***280**, 11 (2000) and references therein.
- [2] An intriguing prospect is the study of a BEC-BCS cross-over, see M. Randeria in *Bose-Einstein condensation*, eds. A. Griffin, D. W. Snoke, and S. Stringari (Cambridge University Press, 1995), and A. J. Leggett, in *Modern Trends in the Theory of Condensed Matter*, eds. A. Pekalski and J. Przystawa (Springer-Verlag, Berlin, 1980), pp. 13–27.
- [3] B. DeMarco, D. S. Jin, *Science*, **285**, 1703 (1999).
- [4] The atom-trap prospect of attaining negative scattering length Cooper-pairing was discussed by H. T. C. Stoof et al., *Phys. Rev. Lett.*, **76**, 10 (1996).
- [5] At least two groups are working towards tuning a a fermi-gas mixture near a Feshbach resonance with the aim of reaching superfluidity: R. Hulet, private communication, and D. Vieira, private communication.
- [6] P. Tommasini et al. cond-mat/9804015; E. Timmermans et al., cond-mat/9805323; *Phys. Rev. Lett.*, **83**, 2691(1999); *Phys. Rep.* **315**, 199 (1999).
- [7] P. D. Drummond et al., *Phys. Rev. Lett.* **81**, 3055 (1998); J. Javanainen, M. Mackie, *Phys. Rev. A*, **59**, R3186-R3189 (1999).
- [8] R. Friedberg and T. D. Lee, *Phys. Rev. B***40**, 6745 (1989).
- [9] Long lived simultaneously trapped atom populations in different spin states have been observed in BECs, see e.g. C. J. Myatt et al., *Phys. Rev. Lett.* **78**, 586 (1997), D. S. Hall et al., *Phys. Rev. Lett.* **81**, 1539 (1998), J. Stenger et al., *Nature (London)* **396**, 345 (1998), and with degenerate fermions, see [3].

- [10] P. G. De Gennes, *Superconductivity of metals and alloys*, W.A. Benjamin, Inc., New York, Amsterdam (1966).
- [11] For convenience we refer to the σ -labels as “spins.”
- [12] This corresponds to a time-dependent variational trial wavefunction that is a product of a molecular BEC wavefunction and a BCS state. For a treatment of the Gaussian trial wavefunction of the BEC, see A. Kerman, P. Tommasini, *Annals of Physics*, **260**, 250 (1997).
- [13] Similarly, second-order perturbation around the off-resonant state gives, to leading order, the expected pressure, $P = (4/5)n\epsilon_F + (4\pi\hbar^2/m)a_{eff}n^2 = (4/5)n\epsilon_F[1 + (5/3\pi)(k_F a_{eff})]$. Although $a_{eff} < 0$ in the $\epsilon > 0$ off-resonant regime, $k_F|a_{eff}| \ll 1$, so that the positive fermi pressure, $(4n\epsilon_F/5)$, dominates.
- [14] In the middle of a spherically symmetric harmonic oscillator trap of frequency ω_T (corresponding to a single particle ground state size $l_T = \sqrt{\hbar/m\omega_T}$), that contains a Fermi-degenerate system of N indistinguishable fermions, the local k_F^{-1} -value equals $k_F^{-1} = \frac{l_T}{2(6N)^{1/3}}$. For typical values, $N \geq 10^5$, this length is much shorter than the extent of the trapped fermion system, $L_T = l_T 2(6N)^{1/3}$.
- [15] Note that $u_k v_k = -\alpha\phi_m/[2E(k)]$ and $F_+ = \Omega^{-1} \sum_{\mathbf{k}} u_k v_k$ diverges linearly as $k \rightarrow \infty$ where $E(k) \propto k^2$. We made the necessary renormalization independent of the cut-off by introducing a molecular excitation energy, ϵ' , calculated in the spirit of second-order perturbation as $\epsilon' = \epsilon - (\alpha^2/2\Omega) \sum_{\mathbf{k}} [\hbar^2 k^2/2m - \epsilon'/2]^{-1}$. Its ultra-violet divergence cancels the F_+ -divergence.
- [16] J. Stenger et al., *Phys. Rev. Lett.*, **82**, 2422 (1999).
- [17] Similar to [15], we renormalised the dynamical equations. We made the renormalization cut-off independent by introducing $\epsilon'(t) = \epsilon(t) - (\alpha^2/2\Omega) \sum_{\mathbf{k}} [\hbar^2 k^2/2m -$

$\epsilon(t)'/2]^{-1}$, a procedure that can be justified in a second-order-perturbation approach provided $|\hbar\dot{\epsilon}| \ll \epsilon^2$.

Figure

Figure 1 Plot showing the fraction of atoms in the molecular condensate (i.e. $|\phi_m|^2/n$), as the fermi/bose system responds to a sudden increase at $t = 1$ of the molecule energy from its initial value, $\epsilon = 8\epsilon_F$ to $\epsilon = 14\epsilon_F$. At $t = 1.2$, ϵ was returned to its initial value. The energy unit, ϵ_F , is the fermi energy of the atoms and the unit of time on the horizontal axis is equal to $\hbar/\epsilon_F$. We choose a realistic atom-molecule coupling constant for which $\alpha\sqrt{n} = \epsilon_F$.

